Question: A plane flies at 10 km./minute in a straight direction and level altitude. When it is directly 5 km. overhead you shoot a heatseeking missile at it, straight up. The missile always points directly at the plane. The missile travels at a constant speed of 11 km./minute. How long does it take to hit the plane?

Hint: The integral of $\sqrt{1 + u^2} du = \ln(u + \sqrt{1 + u^2}) + c$, where c is the constant of integration.

Solution

Assume the shooter is at point (0,0) and the plane is at (0,5) when the missile is fired, traveling easy on the coordinate plane.

Let

x = x-coordinate of missile.

y = y-coordinate of missile.

t = number of minutes since missile was fired.

s = distance traveled by missle.

At all times the slope of the missile = $\frac{dy}{dx} = \frac{5-y}{10t-x}$

Invert that:

 $\frac{dx}{dy} = \frac{10t - x}{5 - y}$

Let's put dy and (5-y) on the same side.

$$(5-y)\frac{dx}{dy} = 10t-x$$

Take the derivative of both sides with respect to y. Use the product rule on the left side.

$$-\frac{dx}{dy} + (5-y)\frac{d^2x}{dy^2} = 10\frac{dt}{dy} - \frac{dx}{dy}$$

$$[1] (5-y)\frac{dx^2}{dy^2} = 10\frac{dt}{dy}$$
Let $\frac{dt}{dy} = \frac{dt}{ds} \circ \frac{ds}{dy}$
 $\frac{ds}{dt} = 11$, so $\frac{dt}{ds} = \frac{1}{11}$

The equation for the arc-length of s is:

$$s = \int \sqrt{1 + \left(\frac{dx}{dy}\right)^2} \, dy$$
$$\frac{ds}{dy} = \sqrt{1 + \left(\frac{dx}{dy}\right)^2}$$

Let's go back to equation [1]:

[1]
$$(5-y)\frac{dx^2}{dy^2} = 10\frac{dt}{dy}$$

$$(5-\gamma)\frac{dx^2}{dy^2} = \frac{10}{11} \times \frac{ds}{dy}$$

$$(5-\gamma)\frac{dx^2}{dy^2} = \frac{10}{11} \times \sqrt{1 + (\frac{dx}{dy})^2}$$
Let $u = \frac{dx}{dy}$

$$(5-\gamma)\frac{du}{dy} = \frac{10}{11} \times \sqrt{1 + u^2}$$

Let's put the expressions with a u on one side.

$$\sqrt{1 + u^2} \, du = \frac{10}{11} \times (5 - y)^{-1} \, dy$$

Using the hint from the integral table:

$$\ln(u + \sqrt{1 + u^2}) = -\frac{10}{11} \times \ln(5 - y) + c$$

Take exp() of both sides:

$$u + \sqrt{1 + u^2} = (5 - y)^{-\frac{10}{11}} \times c$$

Let's solve for c. We know that at t=0 that $\frac{dx}{dy} = 0$. $0 + \sqrt{1 + 0^2} = (5 - 0)^{-\frac{10}{11}} \times c$

$$c = 5^{\frac{10}{11}}$$

$$u + \sqrt{1 + u^{2}} = (5 - y)^{-\frac{10}{11}} \times 5^{\frac{10}{11}}$$
$$u + \sqrt{1 + u^{2}} = (\frac{5 - y}{y})^{-\frac{10}{11}}$$
$$\sqrt{1 + u^{2}} = (\frac{5 - y}{y})^{-\frac{10}{11}} - u$$

Square both sides.

$$1 + u^{2} = \left(\frac{5-y}{y}\right)^{-\frac{20}{11}} - 2u\left(\frac{5-y}{y}\right)^{-\frac{10}{11}} + u^{2}$$
$$1 = \left(\frac{5-y}{y}\right)^{-\frac{20}{11}} - 2u\left(\frac{5-y}{y}\right)^{-\frac{10}{11}}$$
$$2u\left(\frac{5-y}{y}\right)^{-\frac{10}{11}} = \left(\frac{5-y}{y}\right)^{-\frac{20}{11}} - 1$$

Remember that $u = \frac{dx}{dy}$

$$\frac{dx}{dy} = 0.5 \times \left[\left(\frac{5-y}{y} \right)^{-\frac{10}{11}} - \left(\frac{5-y}{y} \right)^{\frac{10}{11}} \right]$$

Take the integral of both sides.

x = 0.5 ×
$$\left[-55 * \left(\frac{5-y}{y}\right)^{\frac{1}{11}} + \frac{55}{21} \left(\frac{5-y}{y}\right)^{\frac{21}{11}}\right] + c$$

Let's solve for c, using the fact that when x=0, y=0.

$$0 = 0.5 \times \left[-55 + \frac{55}{21} \right] + c$$
$$c = \frac{550}{21}$$

So, the equation of the path of the missile is:


$$x = 0.5 \times \left[-55 \times \left(\frac{5-y}{y}\right)^{\frac{1}{11}} + \frac{55}{21} \left(\frac{5-y}{y}\right)^{\frac{21}{11}}\right] + \frac{550}{21}$$

The missile hits the plane when y=5. So solve for x when y=5:

$$x = 0.5 \times [0 + 0] + \frac{550}{21}$$

So, the missile strikes the plane when the plane has traveled 550/21 kilometers. The plane travels at 10 kilometers per minute, so it would have taken 55/21 = 2.6195 minutes = 157.1429 seconds to get that far.

This graph shows the path of the missile. The end at (26.195,5) is where it strikes the plane.



Michael Shackleford -- MathProblems.info